



# V: Type and Breed Characteristics and Uses

Stephen P. Hammack\*

The subject of breeds intrigues most cattle producers. However, for beef cattle production, breeds are only part of a genetic strategy that should include:

- Matching applicable performance or functional levels to environmental, management, and marketing conditions
- Choosing a breeding system, either continuous (in which replacement females are produced within the herd) or terminal (in which replacements are introduced externally)
- Selecting genetic types, breeds, and individuals within breeds that are compatible with the performance level needed and breeding system chosen.

## Genetic classifications and breeds

Cattle can be divided into two basic classifications, *Bos taurus* and *Bos indicus*. *Bos taurus* (non-humped) cattle in the United States originated in the British Isles and western continental Europe. *Bos indicus* (humped, also called Zebu) cattle arose in south central Asia. There are some intermediates containing both *Bos taurus* and *Bos indicus*. Some intermediates created in the United States, particularly in Texas, are commonly referred to as American breeds.

Although it has no strict definition, a breed can be described as animals of common origin with certain distinguishing characteristics that are passed uniformly from parent to offspring. Breed characteristics result from both natural selection and from that imposed by people. Currently, more than 75 breeds can be registered in the United States. For a discussion of breed development, see another publication in this series, E-180, "Texas Adapted Genetic Strategies for Beef Cattle—VI: Creating Breeds and Composites."

Most breeds have registry associations that record ancestry, define and document characteristics, and promote the breed. In this publication, estimates of functional traits or characteristics are based on estimates of breed **averages**, but individuals can vary considerably within breeds.

## Functional traits or characteristics

The major functional traits important in Texas beef production include body size, milking potential, age at puberty, hot climate adaptability, fleshing ability, muscle expression, cutability, and marbling.

### Body size

Body size is best evaluated as weight at a stated level of condition or body fatness. Although there are exceptions, weight tends to be positively related to age—cattle heavier at birth tend to be heavier throughout life. Genetically larger animals usually gain weight faster, and weight and nutritional requirements for body maintenance are

closely related. Size is discussed in Extension publication E-188, "Texas Adapted Genetic Strategies for Beef Cattle—III: Body Size and Milking Level." Frame score, an objective measure of skeletal dimension used to estimate current and future body size, is discussed in Extension publication E-192, "Texas Adapted Genetic Strategies for Beef Cattle—X: Frame Score and Weight."

An important inconsistency occurs in *Bos indicus* in body size relationships. Calves from **dams** containing *Bos indicus* inheritance are relatively small at birth, usually resulting in few difficulties in calving. However, calves by *Bos indicus* **sires** out of *Bos taurus* dams often are relatively large, so calving may be more difficult.

### Milking potential

Milking potential is the genetic capability to produce milk. This is not actual volume of milk produced, which also is influenced by the cow's nutrition and the calf's growth potential and nursing pressure. To characterize milking potential accurately, it should be evaluated relative to body size. Higher-milking females need more nutrients for body maintenance and require higher-quality diets, even when not lactating. Again, consult E-188, "Texas Adapted Genetic Strategies for Beef Cattle—III: Body Size and Milking Level."

### Age at puberty

Age at puberty relates to body size, milking potential, and genetic classification. Smaller cows and higher-milking types usually mature earlier, whereas *Bos indicus* mature relatively late. Although higher-milking females, even large ones, often reach puberty and conceive when relatively young, subsequent fertility can suffer because they may become thin after beginning lactation. While *Bos indicus* types reach puberty relatively late, their productive life usually is longer.

### Hot climate/tropical adaptability

Adaptability to hot climate is highest in cattle with *Bos indicus* inheritance, but some *Bos taurus* are reasonably heat tolerant. Animals with lighter-colored, short hair coats and dark skin are best adapted. High humidity intensifies the effects of heat, especially since hot, humid climates often add the stresses of parasites and low-quality forage. Heat with humidity stresses cattle that fail to shed long, thick hair coats, particularly dark-colored ones. As might be expected, animals tolerant to hot climates are less adapted to cold.

### Fleshing ability

Fleshing ability is the body's capacity to fatten and retain fat. Fleshing ability tends to drop with increases in genetic body size, maintenance requirements relative to size, milking level, and inherent muscularity. Animals poorly adapted to their environment generally are low in fleshing ability. *Bos indicus* often flesh easier than other types on low-quality forage and roughage. Easy-fleshing cattle toler-

\*Professor and Extension Beef Cattle Specialist, The Texas A&M University System.

ate periods of nutritional energy deficiency more easily and, therefore, may reproduce more consistently, but they also over fatten more readily in the feedyard unless properly managed.

### **Muscle expression**

Muscle expression is inherent muscularity, independent of other body tissues. Muscling is the second most important factor in cutability. Heavy-muscled types often are low in fleshing ability, so reproductive efficiency may be reduced.

### **Cutability**

Cutability, or percentage of lean, is usually evaluated in beef carcasses as USDA Yield Grade. Cutability depends on relative amounts of fat (which varies greatly), muscle, and bone (which varies least). When genetic types or breeds are compared for differences in cutability, it is assumed that the breeds have similar nutrition levels. But inherent cutability differences can be readily altered by varying nutrition to achieve similar degrees of fatness.

### **Marbling**

Marbling, or intramuscular fat, is the primary factor determining USDA Quality Grade, an indicator of the palatability factors of tenderness, juiciness, and flavor. Marbling increases with age up to physiological maturity and generally is higher in earlier-maturing and higher-milking types. *Bos indicus* and most heavy-muscled, low-milking types have relatively low marbling. Because marbling relates somewhat to body fatness, especially in comparing breeds or types, there is usually a trade-off between Yield Grade and Quality Grade. As one improves, the other tends to decline. For a discussion of carcass genetic considerations, see E-165, "Texas Adapted Genetic Strategies for Beef Cattle—IX: Selection for Carcass Merit."

An important consideration in determining optimum functional levels is that cattle may differ in performance under varying conditions. For a discussion of this topic, see E-187, "Texas Adapted Genetic Strategies for Beef Cattle—II: Genetic-Environmental Interaction."

## **Breeds and functional types**

Table 1 lists characteristics of the major breeds of cattle in Texas those most numerous or most familiar in the state—by estimates of purebred breed-wide averages. Appendix A lists other breeds with registry associations. Using genetic classification and levels of functional traits, breeds can be grouped into these functional types:

- **British Beef**—British-originated breeds and combinations used for beef production only
- **Continental Beef**—Continental European breeds and derivatives developed exclusively for beef production. These are part of what are sometimes called "exotics."
- **Continental Dual Purpose**—breeds selected for both beef and dairy production in their native areas, mostly Continental Europe, and combinations of beef and dairy breeds. Dual Purpose breeds are used only for beef in the United States. They are the other part of "exotics."
- **Dairy**—originating in western Europe and selected in the United States for dairy purposes only, with beef production as a byproduct
- ***Bos indicus***—of pure or very high-percentage *Bos indicus* background and used only for beef production.
- **American**—includes beef breeds created in the United States from combinations of about one-fourth to one-half tropical-adapted inheritance, typically the *Bos indicus*-derived Brahman, along with British Beef, Continental Beef, or Continental Dual Purpose.

- **Specialty**—includes breeds that cannot be placed logically in any of these groups, often characterized by particular emphasis on certain traits.

## **Matching functional levels to production criteria**

Climate and nutrition are key variables affecting where differing types and breeds can be produced. When cattle are not adapted to climatic conditions, production suffers. In hot, humid climates, cattle benefit from some *Bos indicus* or other tropical-adapted genetics.

Table 2 shows the effects of nutrition on optimum levels of three primary production functions in cow herds. In general, as nutrition declines, the smaller, lower-milking, easier-fleshing types are better adapted and more efficient. This is discussed in detail in E-188, "Texas Adapted Genetic Strategies for Beef Cattle—III: Body Size and Milking Level."

Appropriate functional levels can differ, depending on the breeding system implemented. Cattle for general-purpose, continuous systems must contain a blend of important production traits in both sires and dams.

Conversely, specialized sire and dam types are useful in terminal systems. To reduce cowherd nutritional needs or increase cow stocking rate, terminal dams can be relatively small, complemented by high-growth sires. Maternal ability is unimportant in terminal sires because their heifers are not kept for replacements. For more discussion on breeding systems, see Extension publication E-189, "Texas Adapted Genetic Strategies for Beef Cattle—IV: Breeding Systems."

## **Using functional types**

There is no "best" type or breed for beef production because of extensive variations in climates, production conditions, and market requirements. Characteristics of functional types should be matched to these three factors. Also, breeds extreme in specific functional traits usually should be combined with complementing types in a synergistic crossbreeding plan.

The various functional types have these general uses:

### **British beef**

British beef are widely applicable, with some limits in tropical and subtropical climates. Unlike most types, British Beef breeds can be logically straightbred for commercial production. To take advantage of hybrid vigor, cross them with other breeds of this type and with all other types. British breeds are suitable for general-purpose production as well as both the dam side and sire side of a terminal cross, depending on the target market. These breeds are the foundation of the U.S. beef herd and the basis of comparison for other types in this discussion.

### **Continental beef**

Because of weight-gaining ability and cutability, Continental beef are most effective as terminal sires, especially on smaller cows, particularly those with higher-marbling genetics. However, caution is warranted because they may exhibit potentially heavy birth weights and associated calving problems. If needed, Continental beef influence increases size, muscling, and leanness in females without elevating milk production. In general, do not straightbreed or cross this type with other large cattle. Breeds in this group vary considerably in adaptability to hot climates.

**Table 1. Functional Levels of Major Cattle Breeds in Texas<sup>1</sup>**

Functional Type Breed	Body Size <sup>2</sup>	Milking Potential	Age at Puberty <sup>3</sup>	Hot climate Adaptability	Fleshing Ability	Muscle Expression	Cutability <sup>4</sup>	Marbling
<b>British Beef</b>								
Angus	M	M	H	M	H	M	L	H
Hereford <sup>5</sup>	M	L	M	M	H	M	L	M
Red Angus	M	M	H	M	H	M	L	H
Shorthorn	M	M	M	M	H	M	L	H
<b>Continental Beef</b>								
Charolais	VH	L	L	M	M	VH	VH	M
Chianina	VH	L	L	H	L	H	VH	L
Limousin	M	L	L	M	M	VH	VH	L
<b>Continental Dual Purpose</b>								
Braunvieh	H	H	H	H	M	H	H	M
Gelbvieh	H	H	H	M	M	H	H	M
Maine-Anjou	H	M	M	M	M	H	H	M
Salers	H	M	M	L	M	H	H	M
Simmental	VH	H	M	M	M	H	H	M
<b>Dairy</b>								
Holstein	H	EH	H	M	L	L	M	M
Jersey	L	VH	VH	H	M	VL	VL	VH
<b>Bos Indicus</b>								
Brahman	H	M	VL	EH	H	M	M	L
<b>American</b>								
Beefmaster	H	M	L	VH	H	M	L	L
Braford	M	M	L	VH	H	M	L	L
Brangus	H	M	M	VH	H	M	L	M
Red Brangus	M	M	M	VH	H	M	L	M
Santa Gertrudis	H	M	L	VH	H	M	L	L
Simbrah	H	H	L	VH	M	M	M	L
<b>Specialty</b>								
Texas Longhorn	L	L	M	H	M	L	L	M

<sup>1</sup>Breeds thought to be most numerous or familiar in Texas. Evaluations are estimates of purebred breed-wide averages compiled from research reports, particularly U.S. Meat Animal Research Center, and judgments of Texas A&M Animal Science Department members Ronald Gill, Stephen Hammack, Joe Paschal, James O. Sanders, Todd Thrift, and J. W. Turner. See text for explanation of productive functions, characterized above as: VL=very low, L=low, M=medium, H=high, VH=very high, EH=extremely high. Range exists within these categories, so breeds with the same designation do not necessarily average exactly the same level of trait expression. Also, considerable individual variation exists within breeds.

<sup>2</sup>Evaluated as body weight at the same age and body condition.

<sup>3</sup>VL=very late, L=late, M=medium, H=early, VH=very early

<sup>4</sup>Under similar nutrition. See text for explanation.

<sup>5</sup>Horned and Polled.

**Table 2. Matching Cowherd Functional Levels to Nutrition**

Nutritional Availability <sup>1</sup>	Mature Size	Milking Potential	Fleshing Ability
Low	Low to Medium	Low to Medium	High
Medium	Medium	Medium	Medium to High
High	Medium to High	Medium to High	Medium

<sup>1</sup>Quantity, quality, and consistency of nutrition whether from grazing, harvested forage, or supplemental concentrates.

## Continental Dual Purpose

As with Continental beef, these are best used as terminal sires. Maternal use is appropriate for the major breeds of this type, when nutrition is adequate, to create females that are larger, more muscular, leaner, **and** heavier milking. Use the same cautions as with the Continental beef group in birth weight, straightbreeding, and crossing with other large cattle. Because some of the lesser-known breeds in this group are smaller than the major breeds in the group, they are more applicable as general-purpose than terminal sires.

## Dairy

Dairy for beef production are used primarily to create early-maturing, high-milking females **without** increasing muscle. Dairy influence, particularly the smaller breeds, also should maintain or possibly increase fertility if body condition is maintained. However, it often is difficult to keep dairy breeds and crosses in good flesh on typical rangeland or coarse pasture. Price discounts are common for stocker, feeder, and slaughter animals of visible dairy breeding.

## Bos indicus

*Bos indicus* are used primarily to create hot climate-adapted crossbred females with the highest levels of hybrid vigor and calving ease. Generally, these females are best used in terminal crossing systems. Do not straightbreed or cross these breeds with other cattle containing *Bos indicus* unless climatic conditions are persistently hot and humid.

## American

American breeds are widely applicable, especially for, but not limited to, hot climates. They can be straightbred, crossed with other American breeds, or crossed with all other types except purebred or

high-percentage *Bos indicus*. American breeds can be used effectively in either general-purpose production or in terminal systems. The larger, faster-gaining breeds often are a logical choice for terminal sires used in natural service in hot climates.

## Specialty

Specialty breeds should be used where their particular combinations of unusual traits are needed.

## Summary

Sire choice is greatly influenced by types represented in the cow herd. This subject is discussed in the Extension publication, E-191, "Texas Adapted Genetic Strategies for Beef Cattle—VII: Sire Types for Commercial Herds."

There is much genetic variation within functional types and breeds. However, in creating specific production levels, it is usually more effective to begin by exploiting breed averages of applicable functional types than by searching for genetic outliers in other breeds.

A fundamental challenge in commercial beef production is to match genetic capability with climatic, nutritional, management, and market conditions. Knowing functional levels of types and breeds can help producers optimize animal performance to achieve the highest profit.

## For further reading

To obtain other publications in this Texas Adapted Genetics Strategies for Beef Cattle series, contact your county Extension office or see the Extension Web site <http://tcebookstore.org> and the Texas A&M Animal Science Extension Web site <http://animalscience.tamu.edu>.

---

### Appendix A. Additional Cattle Breeds with Registry Associations

British Beef	Continental Beef			Specialty
Belted Galloway	Belgian Blue	Pinzgauer	Nellore	Ankole Watusi
British White	Blonde d'Aquitaine	RX3	Zebu	Beefalo
BueLingo	Chi Angus	South Devon	<b>American</b>	Corriente
Devon	Marchigiana	Tarentaise	American	Dexter
Galloway	Parthenais	<b>Dairy</b>	Barzona	Geltex
Highland	Piedmontese	Ayrshire	Brahmaine	Salorn
Murray Grey	Romagnola	Brown Swiss	Brahmousin	Senepol
Red Poll	<b>Dual Purpose</b>	Guernsey	Bralers	Texon
Sussex	Amerifax	Milking Shorthorn	Braunbray	Tuli
Welsh Black	Beef Friesian	<b>Bos Indicus</b>	Charbray	Wagyu
White Park	Dutch Belted	Boran	Gelbray	
	Milking Devon	Gir	Hotlander	
	Normande	Indu-Brazil	Santa Cruz	

---

Produced by Agricultural Communications, The Texas A&M University System  
Extension publications can be found on the Web at: <http://tcebookstore.org>

Visit Texas Cooperative Extension at: <http://texasextension.tamu.edu>

*Educational programs of Texas Cooperative Extension are open to all people without regard to race, color, sex, disability, religion, age or national origin.*

Issued in furtherance of Cooperative Extension Work in Agriculture and Home Economics, Acts of Congress of May 8, 1914, as amended, and June 30, 1914, in cooperation with the United States Department of Agriculture. Chester P. Fehlis, Director, Texas Cooperative Extension, The Texas A&M University System.