

# TEXAS FARM POND MANAGEMENT CALENDAR

Billy J. Higginbotham\*

## TEXAS POND MANAGEMENT CALENDAR

JAN	FEB	MAR	APRIL	MAY	JUNE	JULY	AUG	SEPT	OCT	NOV	DEC
LIME										LIME	
			FERTILIZATION								
			AQUATIC WEED CONTROL								
			FARM POND RENOVATION								
		STOCK ADVANCED BASS AND BLUEGILL								STOCK 1"-3" BLUEGILL	
			STOCK 1"-3" BASS								
		STOCK CATFISH								STOCK CATFISH	
			CATFISH SPAWN								
		BASS SPAWN		BLUEGILL SPAWN							
			SUPPLEMENTALLY FEED CATFISH AND/OR BLUEGILL								
			CATFISH - BASS - BLUEGILL HARVEST								
DRAWDOWN											DRAWDOWN
REFILL											

• **LIME** - Agricultural limestone applications are necessary in ponds with a pH below 6.5 and total alkalinity below 20 ppm. Waters in this category may limit fish growth and reproduction. East Texas ponds in particular should be tested before stocking.

Best months: November-January.

• **FERTILIZATION** - Inorganic fertilizer can be utilized to double fish production by increasing the food supply. Use granular 20-20-5 at 100 pounds/surface acre followed by one or two reduced rate applications of 35 pounds/surface acre as needed to maintain the bloom. Liquid fertilizer such as 16-34-0 is also utilized at one

\*Wildlife and Fisheries Specialist, Cooperative Extension Program, Prairie View A&M University.

gallon/surface acre every two weeks in the spring. Never fertilize and lime a pond at the same time. Fertilization should be continued on an annual basis.

**Best months:** Start in early April, end in June.

- **AQUATIC WEED CONTROL** - Initiate chemical weed control when growth begins in the spring. Proper plant identification is essential to providing good control. Do not treat an entire pond with heavy weed infestations in the summer. Spot treat in hot weather to avoid an oxygen depletion.

**Best months:** Start in April or May - spot treat throughout summer as needed.

- **FARM POND RENOVATION** - Ponds that become out of balance due to a stunted fish population and/or undesirable species should be renovated utilizing rotenone. Liquid or powder 5 percent rotenone at 10 pounds powder or one gallon liquid/acre foot of water kills most problematic species. Treated water is safe for livestock. Restocking is possible three weeks after treatment.

**Best months:** April-September (when water temperature is above 70°F).

- **STOCKING** - Use these rates on new or renovated ponds only (no fish present). Muddy ponds and all ponds less than one surface acre should be stocked only with catfish and fathead minnows (at 500/surface acre as supplemental forage). Catfish stocking rates range from 100 to 1000 fingerlings per acre depending on the frequency of feeding. **Best months:** March-May or October-December. Ponds larger than one surface acre are suitable for management of bass-bluegill or bass-bluegill-catfish. Occasionally, supplemental forage species (threadfin shad, fathead minnows, golden shiners and/or redear sunfish are stocked in addition to the bluegill. Stock 20 six to eight inch bass and 30 three inch or larger bluegill per surface acre if available. **Best months:** March-June. If only one to three inch bass and bluegill are available, stock 500 bluegill/surface acre in the fall followed by 50 bass/surface acre the next spring. Catfish stocked with bass-bluegill should be at least as large as the bass fingerlings (stocking rate 100/surface acre). All stocking rates for bass-bluegill-catfish strategies can be doubled if a fertilization program is utilized.

- **SPAWNING** - Catfish stocked alone should not be encouraged to reproduce by adding spawning habitat. However, when stocked with bass and bluegill, the addition of tires, culverts, etc., into the lake is recommended to increase catfish reproduction. Bass initiate spawning in March with some females spawning two or three times before summer. Bluegill are essential as bass forage because of their ability to spawn throughout the summer (May-October), providing abundant forage for maintaining a bass population.

- **SUPPLEMENTAL FEEDING** - Supplemental feeding greatly increases the pounds of catfish that can be supported in ponds and lakes. Bluegill also benefit from a regular feeding program. Use a floating ration containing at least 28 percent crude protein. Feed at three percent body weight or all that will be eaten by the fish in 10-15 minutes. Feed may also be offered once a week during the winter on warm, sunny days only. **Best months:** March-November.

- **HARVEST** - Hook and line harvest is encouraged whenever catfish reach edible size. Catfish produced for income should be removed and sold (if appropriate) by the end of the growing season (November-December). However, since most pondowners simply raise catfish for their own recreation, carry-over of fish from one year to the next is common. Mature fish (3-4 years) may spawn and the resulting offspring stunt or the carrying capacity of the pond (maximum of 1000/pounds/surface acre with daily feeding) will be exceeded if the fish are not routinely harvested. Return all bass less than 15 inches long during the first three years after stocking. From the fourth year on, remove bass caught less than 12 inches, return 12-15 inch bass, and keep or return those bass over 15 inches. Do not remove more than 20 pounds of bass per surface acre per year to prevent overharvest. No limit should be placed on the number of bluegill harvested.

**Best months:** All year.

- **CATCH RECORDS** - Anglers should maintain accurate records on the numbers and sizes of each species caught during each fishing trip. Harvest can then be monitored to determine when restocking and/or restrictions applicable to certain species become necessary.

**Best months:** All year.

- **DRAWDOWN** - Ponds larger than one surface acre containing bass and bluegill may benefit from annual drawdown of one to three feet. This technique concentrates forage (bluegill) and makes them more available to bass. Exposed shoreline areas can be planted with a combination of winter grasses following drawdown to provide nursery habitat for sportfish fingerlings the next spring.

**Best months:** Drawdown in late October, refill before bass spawning season (March).